Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 13 September 2016 (MN LATEX style file v2.2)

On the existence of accretion-driven bursts in massive star formation

D. M.-A. Meyer,[1]∗ E. I. Vorobyov,[2,3] R. Kuiper[1] and W. Kley[1]
[1]*Institut für Astronomie und Astrophysik, Universität Tübingen, Auf der Morgenstelle 10, 72076 Tübingen, Germany*
[2]*Department of Astrophysics, The University of Vienna, Vienna, A-1180, Austria*
[3]*Research Institute of Physics, Southern Federal University, Stachki 194, Rostov-on-Don, 344090, Russia*

arXiv:1609.03402v1 [astro-ph.SR] 12 Sep 2016

Received June 2016; accepted Month day Year

ABSTRACT
Accretion-driven luminosity outbursts are a vivid manifestation of variable mass accretion onto protostars. They are known as the so-called FU Orionis phenomenon in the context of low-mass protostars. More recently, this process has been found in models of primordial star formation. Using numerical radiation hydrodynamics simulations, we stress that present-day forming massive stars also experience variable accretion and show that this process is accompanied by luminous outbursts induced by the episodic accretion of gaseous clumps falling from the circumstellar disk onto the protostar. Consequently, the process of accretion-induced luminous flares is also conceivable in the high-mass regime of star formation and we propose to regard this phenomenon as a general mechanism that can affect protostars regardless of their mass and/or the chemical properties of the parent environment in which they form. In addition to the commonness of accretion-driven outbursts in the star formation machinery, we conjecture that luminous flares from regions hosting forming high-mass star may be an observational implication of the fragmentation of their accretion disks.

Key words: stars: flare – accretion discs stars: protostars – stars: massive

1 INTRODUCTION

Stars form in reservoirs of gas and dust, which collapse under their own gravity. However, a significant fraction of pre-stellar gas, thanks to the conservation of the gas net angular momentum, lands onto a centrifugally balanced circumstellar disc rather than falling directly onto the forming protostar. The manner in which matter accretes from the disc onto the star is still poorly understood and, notably, early spherical collapse models (Larson 1969; Shu 1977) that neglect the disc formation phase cannot explain the variety of mass accretion rates observed in present-day star-forming regions (Vorobyov 2009). In particular, these models yield accretion luminosities that are factors of $10-100$ greater than the mean luminosity measured for nearby star-forming regions.

This so-called "luminosity problem" (Kenyon et al. 1990) can be solved if the accretion history onto the protostar is not smooth or quasi-constant, as predicted by spherical collapse models, but highly time-variable, as it naturally occurs in self-consistent models that follow the transition from clouds to protostellar discs (Dunham & Vorobyov 2012). In these models, protostars spend most of their time in the quiescent phase with low rate of accretion, which is interspersed with short but intense accretion bursts (see Vorobyov & Basu 2006, 2010; Machida et al. 2011; Zhu et al. 2012; Vorobyov & Basu 2015). Spectacular examples of these burst systems are a special class of young stars called FU Orionis objects, which display outbursts of a factor of hundreds in luminosity which last several decades to hundreds of years. Such flares are thought to be due to drastic increases in the mass accretion rate of such young stars (Kley & Lin 1996).

Until recently, it was thought that FU-Orionis-type accretion and luminosity bursts were constrained to occur in the solar mass regime of present-day star formation (Audard et al. 2014). However, recent numerical hydrodynamics simulations of primordial disc formation around the first very massive stars have also revealed the presence of accretion bursts caused by disc gravitational fragmentation followed by rapid migration of the fragments onto the protostar (Stacy et al. 2010; Greif et al. 2012; Smith et al. 2012; Vorobyov et al. 2013; Hosokawa et al. 2016). These studies have revealed highly variable protostellar accretion with multiple bursts, exceeding in numbers their present-day counterparts (DeSouza & Basu 2015). The same process of bursts driven by disc fragmentation operates around primordial super-massive stars, relaxing the ultraviolet photon output and enabling the stellar growth to the limit where general-relativistic instability results in the formation of super-massive black holes (Sakurai et al. 2016).

Consequently, the emerging question is, how universal is variable accretion with episodic bursts in star formation and whether it can be associated to a unique physical mechanism? Time-variability of accretion onto present-day massive stars at the early phase of their formation is a known process (Krumholz et al. 2007; Peters et al. 2010; Kuiper et al. 2011; Klassen et al. 2016). Those studies interpret this phenomenon as a natural consequence of the

∗ E-mail: dominique.meyer@uni-tuebingen.de



Figure 1. Midplane density in the center of the computational domain around the time of the first outburst. (a) The region within 800 AU when a clump forms in a spiral arm $\sim 200\,\mathrm{AU}$ away from the protostar, at a time $18.10\,\mathrm{kyr}$. Panel (b-c) display zooms to illustrate the migration and accretion of a part of the clump at times 18.28, 18.30 and $18.32\,\mathrm{kyr}$, respectively. The density is plotted in g/cm^3 on a logarithmic scale and the size of the panels is in AU.

three-dimensional nature of their self-gravitating numerical simulations, while Kuiper & Yorke (2013) explains it by the interplay between mass accretion, stellar evolution, and radiative feedback. They report how asymmetries can develop in self-gravitating discs and generate an azimuthal anisotropy in the accretion flow onto the protostars. One can particularly notice that in addition to its variable character, it is interspersed with several accretion peaks (see fig. 4 of Klassen et al. 2016). In the above cited references, the sharp increases of the accretion rate are generated in simulations assuming different pre-stellar core masses, ratio of kinetic by gravitational energy β, assuming either a *rigidly* rotating cloud or a turbulent pre-stellar core.

The study of a maser outflow in the high-mass star forming region W75N equally conjectures that "short-lived outflows in massive protostars are probably related to episodic increases in the accretion rates, as observed in low-mass star formation" (Carrasco-González et al. 2015). Additionally, a luminosity outburst of the massive ($\approx 20\,\mathrm{M}_\odot$) young star S255IR-NIRS3 was reported in Fujisawa et al. (2015) by means of 6.7 GHz methanol maser emission. This emission line has been discovered by Menten (1991) and constitutes today a well known-tracer of high-mass star forming regions (see Bartkiewicz et al. 2016, and references therein). Recent observation of the same object show brightness variations that resemble strongly FU-Orionis-type outbursts (Stecklum et al. 2016).

Motivated by the above listed numerical studies and observational arguments, we continue to investigate the burst phenomenon in the high-mass regime of star formation. We follow existing models showing accretion spikes in high-mass star formation, further analyze their nature in the context of the star-disc evolution and conjecture on possible observational implications. This study is organized as follows. In Section 2, we review the methods that we utilise to carry out our high-resolution self-gravity radiation-hydrodynamical simulation of the formation and evolution of a disc surrounding a growing present-day massive protostar generated by the collapse of a *non-rigidly* rotating pre-stellar core. Our outcomes are presented and discussed in Section 3. Particularly, our model also generates such outbursts and we show that they are caused by the rapid migration of disc fragments onto the protostar. Finally, we conclude on their significances in Section 4.

2 NUMERICAL SIMULATION

We perform a 3D numerical radiation hydrodynamics simulation with the PLUTO code (Mignone et al. 2007, 2012) that has been augmented with several physics modules for (i) self-gravity of the gas and (ii) a careful treatment of the proto-stellar irradiation feedback (Kuiper et al. 2010; Kuiper & Klessen 2013). Our model has been carried out using a spherical coordinate system (r,θ,ϕ) mapped with a grid of size $[r_{\rm min}, r_{\rm max}] \times [0, \pi/2] \times [0, 2\pi]$ that is made of $128 \times 21 \times 128$ cells, respectively. The grid resolution expands logarithmically in the radial direction, scales as $\cos(\theta)$ in the polar direction, is uniform in the azimuthal direction and assumes midplane symmetry about $\theta = \pi/2$, such that it allows us to reach sub-AU spatial resolution in the inner region of the disc. The dynamics of the collapse and the accretion flow is calculated in the frame of reference of the fixed protostar that is considered to be evolving inside a static, semi-permeable sink-cell of initial mass $\approx 10^{-3}\,\mathrm{M}_\odot$ with a radius $r_{\rm min} = 10\,\mathrm{AU}$ such that the mass flux through it represents the protostellar accretion rate $\dot{M}$ onto the stellar surface. The creation of other sink cells is not allowed. As described in Kuiper et al. (2010), we utilise $\dot{M}$ to self-consistently update the stellar mass and interpolate the pre-calculated proto-stellar evolutionary tracks of Hosokawa & Omukai (2009) to use them as boundary conditions for the radiation transport module.

We follow the gravitational collapse of a $M_{\rm c} = 100\,\mathrm{M}_\odot$ pre-stellar core of outer radius $r_{\rm max} = 0.1\,\mathrm{pc}$. The radiation is regulated using the dust opacity prescription of Laor & Draine (1993) while the constant gas opacity is taken to be $\kappa_{\rm g} = 0.01\,\mathrm{g\,cm^{-2}}$. This hybrid radiation transport method allows us to carefully treat disc thermodynamics and to generate the dust sublimation front in massive protostellar accretion discs (cf. Vaidya et al. 2009). Our pre-stellar core has the canonical radial density distribution $\rho(r) \propto r^{-3/2}$. We do not consider a rigidly-rotating core but impose a radial angular momentum distribution $\Omega(R) \propto R^{-3/4}$ with $R = r\sin(\theta)$ the cylindrical radius. We impose a ratio of rotational to gravitational energy $\beta = 4\%$, its initial temperature is uniformly taken to $T_{\rm g} = 10\,\mathrm{K}$ and we set its pressure assuming that the gas is at solar metallicity and obeys an ideal equation of state. The dust temperature $T_{\rm d}$ is considered as equal to the gas temperature $T_{\rm g} = T_{\rm d}$. The collapse of the core and the evolution of the accretion disc is followed during $30\,\mathrm{kyr}$.

3 RESULTS

3.1 Episodic accretion of dense gaseous clumps

The midplane density field from the simulation (in $\mathrm{g\,cm^{-3}}$) is shown in Fig. 1 at different evolutionary times (in kyr). While panel (a) represents an overview of the whole disc region (where material orbits the star with approximately Keplerian velocity), panels



Figure 2. Left: accretion rate onto the protostar and mass infall rate onto the disc (in $\mathrm{M}_\odot\ \mathrm{yr}^{-1}$). Right: total luminosity of the protostar (in $\mathrm{L}_\odot$).

(b)-(d) are zooms focusing on the inner region of the disc. The first snapshot (a) shows the circumstellar medium of the protostar at a time $18.10\,\mathrm{kyr}$. In addition to the gas that is being accreted in the innermost $50\,\mathrm{AU}$ of the disc, it develops inhomogeneities by gravitational instability that take the form of spiral arms terminated by dense blobs of gas orbiting at radii $450-800\,\mathrm{AU}$ from the protostar. One of those Truelove-resolved arms ($\lambda/\Delta \sim 6-8$, where λ is the Jeans length and Δ is the grid cell size, see Truelove et al. 1998) experiences a local increase in density and develops an overdense clump of circumstellar material at a radial distance of $\approx 200\,\mathrm{AU}$ to the central star. This is how fragmentation occurs at those radii ($\geqslant 150\,\mathrm{AU}$) predicted by the analytic study of Kratter & Matzner (2006), despite the fact that only marginal fragmentation for discs like the one around our $\approx 4.5\,\mathrm{M}_\odot$ protostar is foreseen (see also Fig. 3b).

By means of angular momentum transfer to the closest spiral arm, the clump migrates towards the protostar. At the onset of migration (Fig. 1a), the closest distance to which the clump can fall, assuming angular momentum conservation, is $\sim 2000\,\mathrm{R}_\odot$, and it decreases to $\sim 260\,\mathrm{R}_\odot$ (Fig. 1b) before to become comparable to the stellar radius of $\sim 90\,\mathrm{R}_\odot$ shortly before entering the sink cell. This value is certainly an upper limit because the angular momentum of the clump decreases during its inward migration thanks to the gravitational interaction with the closest spiral arm (the gravitational torque acting on the clump is negative), as was shown in application to low-mass star formation by Vorobyov & Basu (2006). As a consequence, we expect the clump to fall directly onto the star. However, if the gas temperature in the clump interior exceeds 2000 K, the clump will further contract owing to dissociation of molecular hydrogen. In this case, not captured by our numerical simulations due to limited numerical resolution, its subsequent evolution will depend on the amount of rotational energy in the clump. By extrapolation of the clump properties formed in discs around solar-mass stars showing that about half of the clumps are rotationally supported (see Vorobyov 2016), we expect that part of the clump material will retain in the form of a disc/envelope around the newly formed protostar. This material can still be lost onto the central protostar via the Roche lobe creating an accretion burst though of lesser amplitude, as was shown in Nayakshin & Lodato (2012). Furthermore, high-resolution studies of low-mass and primordial star formation showed that further fragmentation may happen within in the innermost disk regions occupied by our sink cell, likely triggered by already existing outer fragments or efficient H_2 cooling and collisionally induced emission (see Meru 2015; Greif et al. 2012). If fragmentation in the inner disk occurs also in the case of high-mass star formation, this may provoke the formation of a pattern of very small clumps which complex mutual gravitational interactions can make them merging together, orbiting onto the protostar or being dynamically ejected away. This would modify the physics of accretion and the accretion luminosity of the protostar, however, heavy clumps migrating very rapidly will still fall directly onto the stellar surface. To follow this process in detail, however, a much smaller (than 10 AU) sink cell and, as a consequence, much longer integration times are needed. We leave this investigation for a subsequent study.

At time $18.28\,\mathrm{kyr}$, the clump is at about $60\,\mathrm{AU}$ from the protostar (Fig. 1b) and its density reaches $\geqslant 10^{-9}\,\mathrm{g\,cm}^{-3}$. The self-gravitating clump adopts a slightly elongated shape due to the gravitational influence of the massive protostar. At time $18.30\,\mathrm{kyr}$ (Fig. 1c), the stretched clump starts wrapping around the sink cell. A part of its material is accreted onto the protostar, and, consequently, it affects the star's accretion luminosity $L_{\rm acc} \propto GM_\star \dot{M}/R_\star$ (see Section 3.2). When the clump starts migrating towards the star its radius is 20 AU. As the clump approaches the sink cell, its mass has decreased to $\simeq 0.7\,\mathrm{M}_\odot$ and its Hill radius decreases from approximately 35 AU to $\lesssim 10$ AU, causing the clump to lose part of its mass in the diffuse outer envelope. Finally, about $0.55\ \mathrm{M}_\odot$ is accreted through the sink cell. After time $18.32\,\mathrm{kyr}$ (Fig. 1d), the structure of the density field around the sink cell is that of a spiral of unaccreted dense gas material.

This scenario is episodic in the disc evolution and repeats itself at a time about $21.00\,\mathrm{kyr}$ when another clump is similarly accreted. The time interval between recurrent accretion bursts can be determined as the time needed to replenish the disc material lost during the last accretion event $t_{\rm repl} = M_{\rm cl}/\dot{M}_{\rm infall}$, where $M_{\rm cl}$ is the clump mass and $\dot{M}_{\rm infall}$ the mass infall rate onto the disc from the collapsing envelope (see Section 3.2). For $M_{\rm cl} \sim 0.55\ \mathrm{M}_\odot$ and $\dot{M}_{\rm infall} = (2-3)\times 10^{-4}\ \mathrm{M}_\odot\ \mathrm{yr}^{-1}$, typical values for the time of the bursts, the characteristic time of recurrent outbursts is $t_{\rm repl} \approx 2-2.5$ kyr. If $t_{\rm repl}$ is regarded as the time required to increase the disc mass to the limit when the disc fragments, i.e., as the fragmentation timescale, then the clump migration timescale is much shorter than the fragmentation timescale $t_{\rm repl}$. This is indeed evident from Figure 1.

3.2 Accretion-induced strong outbursts

The accretion rate $\dot{M}$ onto the protostar exhibits rapid variations after the disc formation at about $4\,\mathrm{kyr}$ (see Fig. 2a). Comparing $\dot{M}$ measured at $r_{\rm min} = 10$ AU with the smooth infall rate $\dot{M}_{\rm infall}$ from the envelope onto the disc measured at $r = 3000\,\mathrm{AU}$, it becomes evident that the oscillations of $\dot{M}$ from about 10^{-6} to about $10^{-3}\,\mathrm{M_\odot\,yr^{-1}}$ are caused by the disc gravitational instability when filaments and spiral arms of different density and length converge towards the protostar. This correlation between the strength of disc gravitational instability and the protostellar accretion variability is well known in low-mass star formation (Dunham & Vorobyov 2012; Vorobyov & Basu 2015). The envelope, on the other hand, has a smooth density and velocity structure, and its infall rate that gradually increases with time for the initial free-fall time of the mass reservoir.. Note also that the self-similar solution for the a singular isothermal sphere (Shu 1977) computed as the mean isothermal infall rate at $r = 3000\,\mathrm{AU}$, underestimates the accretion onto the protostar by about an order of magnitude (Fig. 2a), as previously noticed in the context of low-mass and high-mass star formation, see Vorobyov (2009) and Banerjee & Pudritz (2007), respectively.

At times 18.29 kyr and $21.30\,\mathrm{kyr}$ the accretion of parts of the heaviest clumps occurs. The total luminosity of the protostar, i.e., the sum of its photospheric luminosity $L_\star$ and its accretion luminosity $L_{\rm acc}$ rises simultaneously (see Fig. 2b). The first luminosity outburst is particularly strong reaching in magnitude $5 \times 10^5\ L_\odot$. The second burst is notably weaker because the intrinsic luminosity of the protostar has greatly increased after the first one. These accretion and luminosity outbursts are a direct consequence of the accretion of in-spiraling clumps that fall onto the protostar because of the loss of angular momentum that has been exchanged with other structures of the disc such as spiral arms, filaments, and arcs. In Fig. 3 we present various disc and stellar properties around the time of the first burst. The accretion rate exhibits a rapid increase from $\dot{M} \sim 10^{-3}\ M_\odot\ \mathrm{yr^{-1}}$ to $\dot{M} \approx 10^{-1}\,\mathrm{M_\odot\,yr^{-1}}$ over a time interval of $\approx 10\,\mathrm{yr}$, while the protostar experiences a flare reaching $\geqslant 5 \times 10^5\,\mathrm{L_\odot}$ (Fig. 3a). As the protostar accretes the clump, its mass increases while the disc mass decreases by the amount of mass of the accreted clump. This results in a remarkable step-like decrease of the disc-to-star mass ratio (see Fig. 3b), making the disc temporarily stable to gravitational fragmentation. It takes a few several kyr for the disc to accrete additional mass from the collapsing envelope and produce another similar fragmentation episode. This phenomenon repeats for as long as there is enough material in the envelope to replenish the disc mass, as was previously found for low-mass (Vorobyov 2010) and primordial star formation (Vorobyov et al. 2013; Hosokawa et al. 2016).



Figure 3. Upper panel: As in Fig. 2 during the time interval corresponding to the first accretion-induced outburst. Bottom panel: Protostellar mass ($\mathrm{M_\odot}$) and disc-to-star mass ratio, for a time interval excluding the initial gravitational collapse where no disc has formed yet. The magenta dots mark the times of the zooms in Fig. 1b-d.

4 DISCUSSION AND CONCLUSION

We have confirmed the presence of strong peaks in the variable accretion history of high-mass protostars, already present in the literature for a wide range of initial conditions of the pre-stellar core. All models have a standard initial density distribution $\propto r^{-3/2}$ that is assumed in the absence of corresponding observations. All rotating models preceding our study assumed cores in sold-body rotation. Turbulent radiation-hydrodynamics simulations include (Krumholz et al. 2007; Peters et al. 2010; Seifried et al. 2011), other studies neglected turbulence. Initial conditions of $M_{\rm c}$ and β-ratio are $100{-}200\,\mathrm{M_\odot}$ with 2% (Krumholz et al. 2007), $100\,\mathrm{M_\odot}$ with 2% (Krumholz et al. 2009; Kuiper et al. 2011), $1000\,\mathrm{M_\odot}$ with 5% (Peters et al. 2010), $100\,\mathrm{M_\odot}$ with $4{-}20\%$ (Peters et al. 2010) and $100; 200\,\mathrm{M_\odot}$ with $10.5; 5.3\%$ (Klassen et al. 2016), respectively, while we use $100\,\mathrm{M_\odot}$ with 4%. Our higher spatial resolution in the inner disc allows us to explicitly capture the fall of forming circumstellar clumps rather than modelling overdense filaments in the disc that wrap onto the star. It enables to study this process, well-known in the low-mass and primordial regime of star formation as the so-called episodic accretion-driven outbursts, which we conclude to be also present in the high-mass regime of contemporary star formation. To confirm this result, we have performed preliminary simulations varying the initial rotation curves of the pre-stellar core. In the subsequent studies, numerical simulations with a smaller sink cell are needed to determine more in detail the final fate of the accreted clumps. However, we expect that the protostellar accretion history remains quantitatively similar in models with sink radii varied by a factor of two, as was earlier shown for low-mass star formation (see Vorobyov & Basu 2015, and references therein).

We obtain the protostellar accretion history which is highly time-variable and shows sudden accretion spikes, as was found in several previous models of massive-star-forming pre-stellar cores and interpreted as caused by azimuthal asymmetries in the accretion flow. The higher spatial resolution of our model reveals that, in addition to variable accretion indeed caused by the asymmetric character of its disc, the growing massive protostar can accrete material of gaseous clumps formed in spiral arms owing to disc grav-

itational fragmentation, which rapidly migrate towards the protostar and induce luminous protostellar outbursts. A similar rapid migration of dense clumps is present in gravitationally unstable discs around solar-mass stars, in primordial discs around the first stars and when high-mass planets form (Baruteau et al. 2011; Vorobyov 2013). Our work indicates that this also applies to the high-mass regime of star formation, which implies a change in the paradigm which, to the best of our knowledge, considered episodic accretion-induced protostellar outbursts as inherent to low-mass and primordial star formation only. Our study shows that it concerns star formation in general, for a wide range of the physical properties such as the initial mass, β-ratio, angular momentum distribution or chemical composition of the parent environment in which stars form.

It supports the consideration of star formation as a process ruled by a common set of mechanisms leading to circumprotostellar structures similarly organized, but scaled-up with respect to each other as a function of the initial properties of their parental pre-stellar cores, as observationally suggested (Shepherd & Churchwell 1996; Fuente et al. 2001; Testi 2003; Keto & Zhang 2010; Johnston et al. 2015). Finally, we propose to consider flares from high-mass protostellar objects as a possible tracer of the fragmentation of their accretion discs. This may apply to the young star S255IR-NIRS3 that has recently been associated to a 6.7 GHz methanol maser outburst (Fujisawa et al. 2015; Stecklum et al. 2016), but also to the other regions of high-mass star formation from which originated similar flares (Menten 1991) and which are showing evidences of accretion flow associated to massive protostars, see e.g. in W3(OH) (Hirsch et al. 2012), W51 (Keto & Klaassen 2008; Zapata et al. 2009) and W75 (Carrasco-González et al. 2015).

ACKNOWLEDGEMENTS

We thank the anonymous reviewer for his valuable comments which improved the quality of the paper. This study was conducted within the Emmy Noether research group on "Accretion Flows and Feedback in Realistic Models of Massive Star Formation" funded by the German Research Foundation under grant no. KU 2849/3-1. E.I.V. acknowledges support from the Austrian Science Fund (FWF) under research grant I2549-N27 and RFBR grant 14-02-00719.

REFERENCES

Audard M., Ábrahám P., Dunham M. M., Green J. D., Grosso N., Hamaguchi K., Kastner J. H., Kóspál Á., Lodato G., Romanova M. M., Skinner S. L., Vorobyov E. I., Zhu Z., 2014, Protostars and Planets VI, pp 387–410

Banerjee R., Pudritz R. E., 2007, ApJ, 660, 479

Bartkiewicz A., Szymczak M., van Langevelde H. J., 2016, A&A, 587, A104

Baruteau C., Meru F., Paardekooper S.-J., 2011, MNRAS, 416, 1971

Carrasco-González C., Torrelles J. M., Cantó J., Curiel S., Surcis G., Vlemmings W. H. T., van Langevelde H. J., Goddi C., Anglada G., Kim S.-W., Kim J.-S., Gómez J. F., 2015, Science, 348, 114

DeSouza A. L., Basu S., 2015, MNRAS, 450, 295

Dunham M. M., Vorobyov E. I., 2012, ApJ, 747, 52

Fuente A., Neri R., Martín-Pintado J., Bachiller R., Rodríguez-Franco A., Palla F., 2001, A&A, 366, 873

Fujisawa K., Yonekura Y., Sugiyama K., Horiuchi H., Hayashi T., Hachisuka K., Matsumoto N., Niinuma K., 2015, The Astronomer's Telegram, 8286

Greif T. H., Bromm V., Clark P. C., Glover S. C. O., Smith R. J., Klessen R. S., Yoshida N., Springel V., 2012, MNRAS, 424, 399

Hirsch L., Adams J. D., Herter T. L., Hora J. L., De Buizer J. M., Megeath S. T., Gull G. E., Henderson C. P., Keller L. D., Schoenwald J., Vacca W., 2012, ApJ, 757, 113

Hosokawa T., Hirano S., Kuiper R., Yorke H. W., Omukai K., Yoshida N., 2016, Accepted to Apj, ArXiv e-prints:1510.01407v2

Hosokawa T., Omukai K., 2009, ApJ, 691, 823

Johnston K. G., Robitaille T. P., Beuther H., Linz H., Boley P., Kuiper R., Keto E., Hoare M. G., van Boekel R., 2015, ApJ, 813, L19

Kenyon S. J., Hartmann L. W., Strom K. M., Strom S. E., 1990, AJ, 99, 869

Keto E., Klaassen P., 2008, ApJ, 678, L109

Keto E., Zhang Q., 2010, MNRAS, 406, 102

Klassen M., Pudritz R. E., Kuiper R., Peters T., Banerjee R., 2016, ApJ, 823, 28

Kley W., Lin D. N. C., 1996, ApJ, 461, 933

Kratter K. M., Matzner C. D., 2006, MNRAS, 373, 1563

Krumholz M. R., Klein R. I., McKee C. F., 2007, ApJ, 656, 959

Krumholz M. R., Klein R. I., McKee C. F., Offner S. S. R., Cunningham A. J., 2009, Science, 323, 754

Kuiper R., Klahr H., Beuther H., Henning T., 2010, ApJ, 722, 1556

Kuiper R., Klahr H., Beuther H., Henning T., 2011, ApJ, 732, 20

Kuiper R., Klahr H., Dullemond C., Kley W., Henning T., 2010, A&A, 511, A81

Kuiper R., Klessen R. S., 2013, A&A, 555, A7

Kuiper R., Yorke H. W., 2013, ApJ, 772, 61

Laor A., Draine B. T., 1993, ApJ, 402, 441

Larson R. B., 1969, MNRAS, 145, 271

Machida M. N., Inutsuka S.-i., Matsumoto T., 2011, ApJ, 729, 42

Menten K. M., 1991, ApJ, 380, L75

Meru F., 2015, MNRAS, 454, 2529

Mignone A., Bodo G., Massaglia S., Matsakos T., Tesileanu O., Zanni C., Ferrari A., 2007, ApJS, 170, 228

Mignone A., Zanni C., Tzeferacos P., van Straalen B., Colella P., Bodo G., 2012, ApJS, 198, 7

Nayakshin S., Lodato G., 2012, MNRAS, 426, 70

Peters T., Banerjee R., Klessen R. S., Mac Low M.-M., Galván-Madrid R., Keto E. R., 2010, ApJ, 711, 1017

Sakurai Y., Vorobyov E. I., Hosokawa T., Yoshida N., Omukai K., Yorke H. W., 2016, MNRAS, 459, 1137

Seifried D., Banerjee R., Klessen R. S., Duffin D., Pudritz R. E., 2011, MNRAS, 417, 1054

Shepherd D. S., Churchwell E., 1996, ApJ, 472, 225

Shu F. H., 1977, ApJ, 214, 488

Smith R. J., Hosokawa T., Omukai K., Glover S. C. O., Klessen R. S., 2012, MNRAS, 424, 457

Stacy A., Greif T. H., Bromm V., 2010, MNRAS, 403, 45

Stecklum B., Caratti o Garatti A., Cardenas M. C., Greiner J., Kruehler T., Klose S., Eisloeffel J., 2016, The Astronomer's Telegram, 8732

Testi L., 2003, in De Buizer J. M., van der Bliek N. S., eds, Galactic Star Formation Across the Stellar Mass Spectrum Vol. 287 of Astronomical Society of the Pacific Conference Series, Intermediate Mass Stars (Invited Review). pp 163–173

Truelove J. K., Klein R. I., McKee C. F., Holliman II J. H., Howell L. H., Greenough J. A., Woods D. T., 1998, ApJ, 495, 821

Vaidya B., Fendt C., Beuther H., 2009, ApJ, 702, 567

Vorobyov E. I., 2009, ApJ, 704, 715

Vorobyov E. I., 2010, ApJ, 723, 1294

Vorobyov E. I., 2013, A&A, 552, A129

Vorobyov E. I., 2016, ArXiv e-prints:1603.07761

Vorobyov E. I., Basu S., 2006, ApJ, 650, 956

Vorobyov E. I., Basu S., 2010, ApJ, 714, L133

Vorobyov E. I., Basu S., 2015, ApJ, 805, 115

Vorobyov E. I., DeSouza A. L., Basu S., 2013, ApJ, 768, 131

Zapata L. A., Ho P. T. P., Schilke P., Rodríguez L. F., Menten K., Palau A., Garrod R. T., 2009, ApJ, 698, 1422
Zhu Z., Hartmann L., Nelson R. P., Gammie C. F., 2012, ApJ, 746, 110